

GUARANTEED ANALYSIS

Crude protein not less than	17.0%
Crude fat not less than	13.0%
Crude fiber not more than	7.0%
Ash not more than	8.0%

Product Forms Available* Catalog #

Extruded Particle	0011746
-------------------	---------

*Other Forms Available On Request

INGREDIENTS

Ground Oat Hulls, Acid Casein, Wheat Flour, Sucrose, Porcine Animal Fat Preserved with BHA, Ground Corn, Dried Whey, Fructose, Ground Wheat, Powdered Cellulose, Wheat Germ, Dried Egg Product, Poultry Fat Preserved with Ethoxyquin, Dicalcium Phosphate, Calcium Carbonate, Corn Oil, Magnesium Sulfate, Potassium Sulfate, Potassium Chloride, Salt, L-ascorbyl-2-polyposphate, Fish Oil (LS), Pyridoxine Hydrochloride, Menadione Dimethylpyrimidinol Bisulfite, Calcium Pantothenate, Cholecalciferol, Folic Acid, Vitamin A Acetate, Inositol, Thiamine Mononitrate, Biotin, Nicotinic Acid, Lecithin, Riboflavin, Vitamin B-12 Supplement, Casein, Manganous Oxide, Zinc Oxide, Ferrous Carbonate, Copper Sulfate, Zinc Sulfate, Calcium Iodate, Cobalt Carbonate, Sodium Selenite.

FEEDING DIRECTIONS

TAD Primate Diet should be fed within 270 days of manufacture. The date of product manufacture is found at the bottom of the back of the bag. Primates generally consume about 2 to 4 percent of their body weight in food each day. A guide for consumption would be 100-200G/5 KG body weight per day. Animals should be fed twice daily.

CHEMICAL COMPOSITION ¹**Nutrients²**

Protein, %	17.0	Sulfur, %	0.35
Arginine, %	0.72	Sodium, %	0.30
Cystine, %	0.19	Chloride, %	0.67
Glycine, %	0.35	Fluorine, ppm	23.7
Histidine, %	0.45	Iron, ppm	230
Isoleucine, %	0.93	Zinc, ppm	97
Leucine, %	1.55	Manganese, ppm	105
Lysine, %	1.20	Copper, ppm	13
Methionine, %	0.47	Cobalt, ppm	0.53
Phenylalanine, %	0.86	Iodine, ppm	1.44
Tyrosine, %	0.82	Chromium (added), ppm	0.01
Threonine, %	0.75	Selenium, ppm	0.24
Tryptophan, %	0.21		
Valine, %	1.20		
Serine, %	0.87		
Aspartic Acid, %	1.14		
Glutamic Acid, %	3.75		
Alanine, %	0.55		
Proline, %	1.76		
Taurine, %	0.00		

Fat (ether extract), %**Fat (acid hydrolysis), %**

Cholesterol, ppm	621
Linoleic Acid, %	2.79
Linolenic Acid, %	0.11
Arachidonic Acid, %	0.04
Omega-3 Fatty Acids, %	0.24
Total Saturated Fatty Acids, %	5.40
Total Monounsaturated Fatty Acids, %	6.17

Fiber (crude), %

Neutral Detergent Fiber³, %

Acid Detergent Fiber⁴, %

Nitrogen-Free Extract (by difference), %

Starch, %	13.6
Glucose, %	4.54
Fructose, %	10.04
Sucrose, %	8.80
Lactose, %	4.51

Total Digestible Nutrients, %**Gross Energy, kcal/gm****Physiological Fuel Value⁵, kcal/g****Metabolizable Energy, kcal/g****Minerals**

Ash, %	7.1
Calcium, %	1.00
Phosphorus, %	0.60
Phosphorus (non-phytate), %	0.51
Potassium, %	0.79
Magnesium, %	0.17

Vitamins

Carotene, ppm	0.1
Vitamin K, ppm	3.0
Thiamin, ppm	25.1
Riboflavin, ppm	10.1
Niacin, ppm	100
Pantothenic Acid, ppm	75
Choline Chloride, ppm	468
Folic Acid, ppm	8.0
Pyridoxine, ppm	14.0
Biotin, ppm	0.2
Vitamin B-12, mcg/kg	35
Vitamin A, IU/g	23
Vitamin D-3 (added), IU/g	3.0
Vitamin E, IU/kg	2
Ascorbic Acid, ppm	700

Calories provided by:

Protein, %	18.340
Fat (ether extract), %	36.410
Carbohydrates, %	45.251

*** Product Code**

1. Formulation based on calculated values from the latest ingredient analysis information. Since nutrient composition of natural ingredients varies and some nutrient loss will occur due to manufacturing processes, analysis will differ accordingly.

2. Nutrients expressed as percent of ration except where otherwise indicated. Moisture content is assumed to be 10.0% for the purpose of calculations.

3. NDF = approximately cellulose, hemicellulose and lignin.

4. ADF = approximately cellulose and lignin.

5. Physiological Fuel Value (kcal/gm) - Sum of decimal fractions of protein, fat and carbohydrate (use Nitrogen Free Extract) x 4,9,4 kcal/gm respectively.